

Stimulated generation: extraction of energy from balanced flow by near-inertial waves

Cesar B. Rocha¹†, Gregory L. Wagner² and William R. Young¹

¹Scripps Institution of Oceanography, University of California, San Diego

²Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology

(Received xx; revised xx; accepted xx)

We study stimulated generation—the transfer of energy from balanced flows to existing internal waves—using an asymptotic model that couples barotropic quasi-geostrophic flow and near-inertial waves with e^{imz} vertical structure on the f -plane. A detailed description of the conservation laws of this vertical-plane-wave model illuminates the mechanism of stimulated generation associated with vertical vorticity and lateral strain. There are two sources of wave potential energy, and corresponding sinks of balanced kinetic energy: the refractive convergence of wave action density into anti-cyclones (and divergence from cyclones); and the enhancement of wave-field gradients by geostrophic straining. We quantify these energy transfers and describe the phenomenology of stimulated generation using numerical solutions of an initially uniform inertial oscillation interacting with mature freely evolving 2D turbulence. In all solutions, stimulated generation co-exists with a transfer of balanced kinetic energy to large scales via vortex merging. And geostrophic straining accounts for most of the generation of wave potential energy, representing a sink of 10-20% of the initial balanced kinetic energy. But refraction is fundamental because it creates the initial eddy-scale lateral gradients in the near-inertial field that are then enhanced by advection. In these quasi-inviscid solutions, wave dispersion is the only mechanism that upsets stimulated generation: with a barotropic balanced flow, lateral straining enhances the wave group velocity, so that waves accelerate and rapidly escape from straining regions. This wave escape prevents wave energy from cascading to dissipative scales.

Key words:

1. Introduction

The inverse cascade, acting on balanced ocean macroturbulence, transfers energy towards large spatial scales. But a statistically steady ocean circulation requires energy dissipation at the same rate as it is supplied by wind. Thus equilibration of the ocean macroturbulence requires ageostrophic processes, acting in opposition to the inverse cascade, to produce a transfer of energy towards the centimeter scales at which molecular viscosity is effective. Mechanisms that might result in this down-scale transfer include, but are not limited to, surface and benthic boundary-layer turbulence, lee-wave generation by mesoscale eddies negotiating bottom topography, and the spontaneous generation of

† Email address for correspondence: crocha@ucsd.edu

TABLE 1. Summary of model-based studies of energy extraction from balanced flow by near-inertial waves.

Study	Framework	Energy transfers from a mean flow to <i>existing</i> near-inertial waves are referred to as
Gertz & Straub (2009)	Barotropic 2D double-gyre solutions coupled with forced 3D near-inertial waves.	<i>2D-to-3D energy transfer</i>
Thomas (2012)	Near-inertial waves in a baroclinic geostrophic flow undergoing frontogenesis.	<i>deformation shear production</i>
Taylor & Straub (2016)	Boussinesq channel-flow with both high- and low-frequency forcing.	<i>advective sink</i>
Barkan <i>et al.</i> (2016)	Boussinesq channel-flow with both high- and low-frequency forcing.	<i>direct extraction</i>
Shakespeare & Hogg (2017)	Boussinesq channel-flow with low-frequency forcing. Spontaneous generation in the surface layer and stimulated generation in the interior.	<i>interior amplification</i>

internal waves by upper-ocean frontal instabilities; see Nagai *et al.* (2015) for a recent review.

We focus on a mechanism first identified by Gertz & Straub (2009): externally-forced near-inertial waves might provide an energy sink for large-scale balanced flow. Since Gertz & Straub (2009), several other studies, summarized in table 1, have argued for significant energy transfer from balanced flows to near-inertial waves. A common aspect of the studies in table 1 is that the near-inertial waves are first introduced by external forcing (e.g., wind) at the inertial frequency and then grow by extracting energy from the balanced flow.

(An exception in table 1 is the study of Shakespeare & Hogg (2017), in which near-inertial waves are generated spontaneously—without external forcing—at density fronts near the surface. These waves then radiate vertically downwards into the interior and amplify by extracting energy from deep balanced flow. We have included Shakespeare & Hogg (2017) in table 1 because, as far as deep interior amplification is concerned, the details of the shallow generation process are probably immaterial.)

The studies in table 1 have shown that externally generated near-inertial waves can extract energy from a pre-existing balanced flow. Those studies, however, have diverse methodology and diagnostic framework, so there is not a consensus that the observed amplification of near-inertial waves results from a single mechanism. In other words, it is possible that near-inertial wave amplification occurs through a variety of mechanisms and each instance must be analyzed and understood on its own peculiar terms. But in a certain limit, described in section 2.2, there is a single underlying mechanism—stimulated generation—that is responsible for energy transfer between waves and balanced flow (Xie

& Vanneste 2015, XV hereafter). While it is unclear whether this limit applies to the studies in table 1, XV provides a robust yet simple theoretical framework for studying energy transfers between balanced flows and near-inertial waves.

Using a variational formulation of the generalized Lagrangian-mean, XV derived a phase-averaged model of the coupling between weakly non-linear near-inertial waves and quasi-geostrophic (QG) flow. Wagner & Young (2016) derived a similar coupled model via Eulerian multiple-time expansion; these authors include the second harmonic of the primary near-inertial wave and simplify the wave dynamics by assuming moderate QG vertical shears. In both coupled models the near-inertial waves are governed by the equation of Young & Ben Jelloul (1997) (YBJ hereafter) and the balanced flow satisfies QG dynamics—the waves contribute phase-averaged quadratic terms at the same order of the QG potential vorticity (PV). Salmon (2016) provides a useful perspective on this “NIW-QG” model; without assuming that the waves are near-inertial, and within a single variational framework, Salmon unifies XV’s model with the wave-mean flow models of Bühler & McIntyre (1998) and Wagner & Young (2015). Salmon also emphasizes a revealing analogy between vortex-wave interactions and classical electrodynamics.

To distinguish energy extraction by existing waves from *spontaneous generation*, and to complete an electrodynamic analogy, XV refer to the transfer of energy from balanced flow to externally forced near-inertial waves as *stimulated generation*. The more widely studied process of spontaneous generation is the emission of internal waves arising from the slow evolution of balanced flow in the absence of external forcing at wave frequencies (Vanneste 2013). Spontaneous generation is inefficient at small and moderate Rossby numbers and its global impact on ocean energetics is probably small (Danioux *et al.* 2012; Nagai *et al.* 2015). And spontaneous generation is localized at sharp submesoscale fronts with order-one Rossby number (e.g., Shakespeare & Hogg 2017) while the stimulated variety operates even at small Rossby numbers characteristic of most interior oceanic conditions, provided only that internal waves are introduced by external forcing. Throughout the ocean, internal waves are reliably forced by wind and tides and thus stimulated generation is a leading contender as a mesoscale energy sink.

Although XV and Wagner & Young (2016) use significantly different approaches, their results are consistent with one another. This consistency indicates that the NIW-QG model provides the unique *small-amplitude* evolution equations describing the interaction between near-inertial waves and geostrophic flow. In the small-amplitude limit the flow can be unambiguously separated into weakly non-linear internal waves and quasigeostrophic eddies, with perturbative coupling between waves and eddies (Salmon 2016). To the extent that the studies in table 1 are also in this weak-interaction limit, their results should—in principle—be described the NIW-QG model. “In principle” because the distinction between Lagrangian-mean and Eulerian-mean velocities complicates the diagnosis of energy transfers between eddies and waves (see section 6 for further discussion) and because the Rossby number is large in some studies (e.g., Barkan *et al.* 2016). Frontal sharpening and low Richardson number processes, which are described in Thomas (2012), are outside the scope of the NIW-QG model.

XV emphasize that a central feature of the NIW-QG model is that there are two integral energy conservation laws for: (1) near-inertial kinetic energy and (2) the sum of near-inertial potential energy and total balanced energy. The inevitable reduction of near-inertial length scales by advection and refraction is accompanied by an increase in wave potential energy and, because of conservation law (2), a reduction in the energy of the balanced flow. These features, and the necessity of an externally forced wave, are the defining characteristics of stimulated generation.

Here we investigate perhaps the simplest example of stimulated generation obtained

from the NIW-QG model by assuming barotropic QG flow and vertically-planar near-inertial waves. Because the balanced flow is barotropic, while the near-inertial wave is three dimensional, this “vertical-plane-wave model” resembles the original scenario of [Gertz & Straub \(2009\)](#). We show that the convergence of wave kinetic energy into anticyclones and geostrophic straining of the waves reduces the wave length scale, amplifies gradients of wave amplitude, and converts balanced kinetic energy into near-inertial potential energy.

2. The vertical-plane-wave model

The vertical-plane-wave model is obtained by assuming barotropic balanced flow, with streamfunction $\psi(x, y, t)$, a uniform background buoyancy frequency N_0 , and a single vertically propagating wave with vertical structure e^{imz} and back-rotated wave velocity $\phi(x, y, t)$. With these idealizations, appendix A derives the vertical-plane-wave model starting from the phase-averaged equations of [Wagner & Young \(2016\)](#); XV obtain the same model from their version of the phase-averaged equations. In either case, the leading-order wave plus the leading-order balanced velocity (u, v, w) , pressure p , and buoyancy b are:

$$u + iv = e^{i\varpi} \phi - \psi_y + i\psi_x; \quad (2.1)$$

$$w = im^{-1} e^{i\varpi} \partial\phi + cc; \quad (2.2)$$

$$p = -i\eta e^{i\varpi} \partial\phi + cc + f_0\psi; \quad (2.3)$$

$$b = m\eta e^{i\varpi} \partial\phi + cc. \quad (2.4)$$

Above, $\varpi = mz - f_0t$ is the phase of the vertical plane wave, $\eta = f_0\lambda^2$ is the wave “dispersivity,” where $\lambda = N_0/f_0 m$ is a horizontal scale, cc denotes complex conjugate, and

$$\partial \stackrel{\text{def}}{=} \frac{1}{2}(\partial_x - i\partial_y) \quad (2.5)$$

is a differential operator. The complex field $\phi(x, y, t)$ in (2.1) is the back-rotated velocity of the near-inertial waves; in (2.2)–(2.4) the other wave fields are expressed in terms of $\partial\phi$. The compact representation of the wave variables in terms of ϕ follows YBJ.

The balanced variables are represented by the streamfunction ψ . Because the Stokes pressure correction is negligible for near-inertial waves ([Wagner & Young 2016](#)), the eddies are balanced in the sense that $\psi = \bar{p}/f_0$, where \bar{p} is either the Eulerian-mean or Lagrangian-mean pressure. Moreover, the velocities obtained from ψ in (2.1) are Lagrangian-mean velocities, e.g., the velocity $(-\psi_y, \psi_x)$ advects the material invariant potential vorticity in (2.7) below. We have lightened the notation by using ψ , rather than ψ^L ; the implicit L is particularly important in section 6.

The PV of the balanced flow is expressed in terms of ψ and ϕ by

$$q = \underbrace{\Delta\psi}_{\stackrel{\text{def}}{=} \zeta} + \frac{1}{f_0} \underbrace{\left[\frac{1}{4}\Delta|\phi|^2 + \frac{i}{2}J(\phi^*, \phi) \right]}_{\stackrel{\text{def}}{=} q^w}, \quad (2.6)$$

where $\Delta \stackrel{\text{def}}{=} \partial_x^2 + \partial_y^2$ is the horizontal Laplacian and $J(f, g) \stackrel{\text{def}}{=} f_x g_y - f_y g_x$ is the Jacobian, and the superscript star $*$ denotes complex conjugation. Equation (2.6) is the “inversion relation”: q and ϕ determine the Lagrangian-mean flow via $\psi = \Delta^{-1}(q - q^w)$ where q^w defined in (2.6) is the “wave potential vorticity.” Once ψ is obtained by inversion, the field equations (2.7) and (2.8) below can be time-stepped.

Using the generalized Lagrangian-mean formulation [Bühler & McIntyre \(1998\)](#) showed

that the assumption of weak interaction between internal waves and balanced flow results in wave-averaged term q^w contributing to the materially conserved PV; see also [Grimshaw \(1975\)](#). In (2.6) the wave-averaged feedback on the balanced flow is expressed concisely in terms of the backrotated velocity ϕ via the quadratic terms in q^w .

2.1. The evolution equations

The balanced flow evolves according to PV advection

$$q_t + J(\psi, q) = D_q; \quad (2.7)$$

the back-rotated velocity satisfies the wave equation

$$\phi_t + J(\psi, \phi) + \phi \frac{1}{2} \zeta - \frac{1}{2} \eta \Delta \phi = D_\phi. \quad (2.8)$$

D_q and D_ϕ in (2.7) and (2.8) are dissipative terms described below.

The wave equation (2.8) is the YBJ model in the case where the near-inertial wave has e^{imz} structure. The back-rotated wave velocity, ϕ , evolves through dispersion—the last term on the left of (2.8)—and nonlinear advection and refraction by the second and third terms in (2.8). Without advection, (2.8) is analogous to Schrödinger’s equation (e.g., [Landau & Lifshitz 2013](#), page 51). The relative vorticity, $\zeta = \Delta\psi$, is the potential, with negative ζ a well, and the “dispersivity,” $f_0\lambda^2$, is Planck’s constant ([Balmforth et al. 1998](#); [Balmforth & Young 1999](#); [Danioux et al. 2015](#)). The quantum analogy may be useful for some readers, but it is not necessary for the understanding of the results below.

The terms on the right of (2.7) and (2.8), D_q and D_ϕ , represent small-scale dissipation. Small-scale dissipation is necessary to absorb the forward transfers of potential enstrophy and wave kinetic and potential energies in the numerical solutions reported below. We find that biharmonic diffusion and viscosity,

$$D_q = -\kappa_e \Delta^2 q \quad \text{and} \quad D_\phi = -\nu_\phi \Delta^2 \phi, \quad (2.9)$$

are sufficient to extend the spectral resolution compared to Laplacian dissipation. In practice, we choose κ_e and ν_ϕ so that the highest 35% of the modes lie in the dissipation range and aliased wavenumbers are strongly damped.

2.2. The small-amplitude limit and the validity of the NIW-QG approximation

The development of the NIW-QG model is ordered first by assuming that the waves are weak in the sense that

$$\epsilon \stackrel{\text{def}}{=} \frac{U_w}{f_0 L} \ll 1. \quad (2.10)$$

Above, L is a characteristic scale of both waves and balanced flow and U_w is a characteristic near-inertial wave velocity. The other small parameter in the expansion is the Rossby number of the balanced flow

$$Ro \stackrel{\text{def}}{=} \frac{U_e}{f_0 L} \ll 1, \quad (2.11)$$

where U_e is the eddy velocity. The inequalities in (2.10) and (2.11) must be satisfied in order to obtain the NIW-QG system. But [XV](#) and [Wagner & Young \(2016\)](#) make a third assumption: $Ro = \epsilon^2$, or equivalently that $U_e = \epsilon U_w$. The resulting distinguished limit,

$$\epsilon \rightarrow 0, \quad \text{with} \quad Ro = \epsilon^2, \quad (2.12)$$

promotes the importance of wave-averaged effects so that q^w appears at an early, and accessible, order in the expansion. Thus (2.12) is for convenience rather than necessity.

TABLE 2. Description of parameters of Lamb-Chaplygin simulation. The initial condition have Rossby number $Ro = U_e k_e / f_0 \approx 0.05$, wave dispersivity $\hbar = f_0 \lambda^2 k_e / U_e \approx 1$, and wave amplitude $\alpha = Ro(U_w/U_e)^2 \approx 3.75$.

Parameter	Description	Value
$R = 2\pi k_e^{-1}$	Dipole radius	$L_d/15 \approx 84$ km
U_e	Dipole strength	5×10^{-2} m s $^{-1}$
U_w	NIW speed	5×10^{-1} m s $^{-1}$
N_0	Buoyancy frequency	5×10^{-3} s $^{-1}$
f_0	Coriolis frequency	10^{-4} s $^{-1}$
$2\pi m^{-1}$	NIW vertical wavelength	325 m
κ_e	PVbiharmonic diffusivity	5×10^7 m 4 s $^{-1}$
ν_w	NIW biharmonic viscosity	1×10^7 m 4 s $^{-1}$
N	Number of modes	512
L	Domain size	$2\pi \times 200$ km

The asymptotic ordering in (2.12) does not imply that the QG–NIW system fails for weaker waves, i.e., if U_w is comparable to, or even smaller than, U_e . Making U_w weaker than U_e delays wave-averaged effects to longer times—it does not, *per se*, invalidate the expansion. The main problem with the weak-wave limit is that other physics, not considered in the NIW–QG system, will contend with wave-averaged effects on ultra-long time scales. For example, even without waves, order Ro^2 ageostrophic effects modify the evolution of balanced flow and produce departures from QG (e.g., see Muraki, Snyder & Rotunno 1999).

To summarize: the main conditions for the validity of the QG–NIW system are (2.10) and (2.11); additionally, validity of the wave equation (2.8) requires that the wave frequency is close of f_0 . The weak-wave limit $U_w/U_e \rightarrow 0$ is valid within the NIW–QG framework: in that limit the system reduces to the barotropic potential vorticity equation and the YBJ equation for a passive wave field.

The standard QG approximation is used successfully even when $Ro \sim 1$ (Hoskins 1975) and we expect the NIW–QG model to enjoy similar success if \ll is replaced by $<$ in (2.11). Flows with $Ro > 1$ eddies, such as those reported in Barkan *et al.* (2016), evolve on time scales close to f_0^{-1} , and an Eulerian decomposition into near-inertial waves and eddies is ill-defined unless there is spatial scale separation between eddies and waves. These large Rossby number flows are outside the purview of the NIW–QG model (XV; Wagner & Young 2016).

2.3. An illustrative solution: the Lamb-Chaplygin dipole

As a preamble to our discussion of stimulated generation in freely-evolving 2D turbulence, we consider an example in which the initial QG flow is the Lamb-Chaplygin dipole—see figure 1. This dipole is an exact solution of the Euler equations on an infinite two-dimensional plane where the vorticity is confined to a circle of radius R (Meleshko & Van Heijst 1994). The relative vorticity, steady in a frame moving at uniform zonal velocity U_e , is

$$\zeta = \frac{2U_e\kappa}{J_0(\kappa R)} \begin{cases} J_1(\kappa r) \sin \theta, & \text{if } r \leq R, \\ 0, & \text{if } r \geq R. \end{cases} \quad (2.13)$$




FIGURE 1. Snapshots of the Lamb-Chaplygin dipole solution with parameters presented in table 2. Contours depict potential vorticity, $q/(U_e k_e) = [-1.5, -0.5, 0.5, 1.5]$, with dashed lines showing negative values. The upper row shows the wave action density $|\phi|^2/2f_0$. The lower row shows the wave buoyancy; the buoyancy scale is $B = k_e m U_w f_0 \lambda^2$. These plots only show the central $(1/5)^2$ of the simulation domain.

Above $r^2 = (x - x_c)^2 + (y - y_c)^2$ is the radial distance from the center (x_c, y_c) , $\tan \theta = (y - y_c)/(x - x_c)$, and J_n is the n 'th order Bessel function. The matching condition at $r = R$ is that $J_1(\kappa R) = 0$ and the smallest solution is $\kappa R \approx 3.8317$. If there is no coupling to the wave ϕ , then the dipole (2.13) is a solution of the QG equation (2.7).

We strongly perturb the dipole in (2.13) by seeding a wave with initial velocity:

$$\phi(x, y, t = 0) = \frac{1+i}{\sqrt{2}} U_w. \quad (2.14)$$

If there was no dipole, this initial condition produces a spatially uniform near-inertial oscillation with speed U_w . Further parameters of this solution are summarized in table 2; note $U_w = 10U_e$.

We integrate the vertical plane-wave model using a standard collocation Fourier spectral method. We evaluate the quadratic non-linearities, including in the wave potential vorticity (2.6), in physical space and transform the product into Fourier space. We time march the spectral equations using an exponential time differencing method with a fourth order Runge-Kutta scheme—for details, see Kassam & Trefethen (2005) and Cox & Matthews (2002).

With the initial condition in (2.14), q^w in (2.6) and $\nabla\phi$ are both zero at $t = 0$. The refractive term in the wave equation (2.8), however, immediately imprints itself onto ϕ thus creating non-zero $\nabla\phi$ and non-zero q^w . Once refraction has created gradients in ϕ , the advective term in (2.8) can further distort ϕ and increase $\nabla\phi$. This scale reduction




FIGURE 2. (a) Snapshot of q (contours) and ζ (colors) at $t \times U_e k_e = 20$. (b) Snapshot of q (contours) and wave PV q^w (colors). Both black lines and colors depict the contour levels $[-1.5, -0.5, 0.5, 1.5] \times (U_e k_e)$. Solid lines and reddish colors depict positive values; dashed lines and greenish colors show negative values.

of ϕ is most evident in the wave buoyancy shown in lower row of figure 1. Figure 1 also shows the well-known focussing of waves into the negative vortex. But the wave feedback on the mean flow through q^w then results in distortion and shearing of the dipole so that the negative vortex loses its integrity; the lop-sided dipole then starts to drift. Once the negative vortex is distorted to small scales it no longer acts as an effective potential well: the trapping of wave energy by the deformed anti-cyclone is weaker than in the initial condition. In fact, figure 1, which shows the materially conserved PV q , understates the development of small scales in the relative vorticity ζ : figure 2 shows that both ζ and q^w develop small scales with significant cancellation resulting in the relatively smooth field $q = \zeta + q^w$ shown in figure 1. Thus at the final time in figure 1 the waves are no longer strongly trapped in the region with $\zeta < 0$. This phenomenology, including significant cancellation between ζ and q^w , is also characteristic of wave-modified two-dimensional turbulence in section 4.

To understand the results in figure 1 and quantify the stimulated generation of wave energy, we need to understand the conservation laws of the vertical-plane-wave model.

3. Conservation laws of the vertical-plane-wave model

XV noted that the vertical-plane-wave model in (2.6) through (2.8) inherits two quadratic conservation laws from the parent NIW-QG model: if there is no dissipation then wave action,

$$\mathcal{A} \stackrel{\text{def}}{=} \frac{|\phi|^2}{2f_0}, \quad (3.1)$$

and the energy,

$$\mathcal{E} \stackrel{\text{def}}{=} \frac{1}{2} |\nabla \psi|^2 + \frac{1}{4} \lambda^2 |\nabla \phi|^2, \quad (3.2)$$

are both separately conserved. Following Bretherton & Garrett (1968), the action in (3.1) is the wave energy divided by the intrinsic frequency; YBJ observed that to leading order the wave energy is only kinetic and the intrinsic frequency in (3.1) is the inertial frequency f_0 .

The conserved energy density \mathcal{E} in (3.2) is the sum of the kinetic energy of the balanced

flow

$$\mathcal{K} \stackrel{\text{def}}{=} \frac{1}{2} |\nabla\psi|^2, \quad (3.3)$$

and the potential energy of the near-inertial waves,

$$\mathcal{P} \stackrel{\text{def}}{=} \frac{1}{2} b^2 / N_0^2 = \frac{1}{4} \lambda^2 |\nabla\phi|^2. \quad (3.4)$$

Above, $b \propto \partial\phi$ is the wave buoyancy defined in (2.4). We emphasize again that ψ is the streamfunction of the Lagrangian-mean flow, which is in geostrophic balance; the Eulerian-mean flow is not in balance. Thus we refer to \mathcal{K} is the kinetic energy of the balanced flow.

XV explain the physical basis of stimulated generation by noting that balanced kinetic energy \mathcal{K} can be converted into wave potential energy \mathcal{P} while conserving the integral of the total energy \mathcal{E} in (3.2). Indeed, this conversion *must* occur if $\nabla\phi$ is increased by a combination of refraction and advection in the wave equation (2.8). In the example shown in figure 1, the initial wave field in (2.14) has infinite spatial scale and therefore there is no wave potential energy at $t = 0$. The subsequent evolution in figure 1 involves creation of non-zero $\nabla\phi$, corresponding to gain of \mathcal{P} at the expense of \mathcal{K} : this is stimulated generation of near-inertial waves.

To substantiate this intuition, and diagnose results from our simulation of wave modified two-dimensional turbulence, we develop the conservation laws corresponding to (3.1) and (3.2) in more detail.

3.1. Action conservation equation and action flux

Multiplying the wave equation (2.8) by ϕ^* and adding to the complex conjugate we obtain a conservation equation for action density

$$\partial_t \mathcal{A} + J(\psi, \mathcal{A}) + \nabla \cdot \mathcal{F} = \underbrace{\frac{1}{2f_0} (\phi^* D_\phi + \phi D_{\phi^*})}_{\stackrel{\text{def}}{=} D_{\mathcal{A}}}, \quad (3.5)$$

where the flux of near-inertial wave action is

$$\mathcal{F} \stackrel{\text{def}}{=} \frac{i}{4} \lambda^2 (\phi \nabla \phi^* - \phi^* \nabla \phi). \quad (3.6)$$

The local conservation law (3.5) shows how the wave action \mathcal{A} changes due to geostrophic advection and divergence of the wave flux and dissipation—the second, third, and fourth terms in (3.5).

The wave action flux \mathcal{F} is analogous to the probability current of quantum mechanics (e.g., Landau & Lifshitz 2013, pg. 57). Using the polar representation $\phi = |\phi|e^{i\Theta}$, the wave action flux \mathcal{F} can also be written as

$$\mathcal{F} = \mathcal{A} \eta \nabla \Theta, \quad (3.7)$$

where recall that $\eta = f_0 \lambda^2$ is the dispersivity. In (3.7), $\eta \nabla \Theta$ is the “generalized group velocity” of hydrostatic near-inertial waves, i.e., \mathcal{F} is the generalized group velocity times the action density \mathcal{A} . We use the term “generalized” because no WKB-type scale separation is required to obtain the results above. The connection to standard internal-wave group velocity is quickly verified by considering a plane near-inertial wave with $\Theta = kx + ly$, yielding $\eta \nabla \Theta = N_0^2(k, l) / f_0 m^2$.

Another useful identity involving the action flux \mathcal{F} is

$$\nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) = \frac{i}{2} \lambda^2 J(\phi^*, \phi), \quad (3.8)$$

where $\hat{\mathbf{k}}$ is the unit vector perpendicular to the (x, y) -plane. Using (3.8), and the definition

of action in (3.1), the wave PV q^w in (2.6) can be written as

$$q^w = \frac{1}{2}\Delta\mathcal{A} + \eta^{-1}\nabla\cdot(\hat{\mathbf{k}}\times\mathcal{F}). \quad (3.9)$$

Denoting an average over the domain by $\langle \rangle$, and assuming that the action flux divergence $\nabla\cdot\mathcal{F}$ vanishes after integration, we obtain from (3.5)

$$\frac{d\langle\mathcal{A}\rangle}{dt} = \varepsilon_{\mathcal{A}}. \quad (3.10)$$

where $\varepsilon_{\mathcal{A}} \stackrel{\text{def}}{=} \langle\phi^*D_{\phi} + \phi D_{\phi^*}\rangle/(2f_0)$ is the domain average of the dissipative term on the right of (3.5). In the example shown in Figure 1 the total action $\langle\mathcal{A}\rangle$ is conserved to within 1% over the course of the integration.

3.2. Ehrenfest's theorem

The quantum analogy suggests that we should seek an analog of Ehrenfest's theorem (the quantum equivalent of Newton's law that force equals mass times acceleration). Thus in Appendix B we develop a local conservation law for \mathcal{F} . The domain average of that result is

$$\frac{d\langle\mathcal{F}\rangle}{dt} = \hat{\mathbf{k}}\times\langle\zeta\mathcal{F}\rangle - \hat{\mathbf{k}}\times\langle(\mathcal{F}\cdot\nabla)\nabla\psi\rangle - \eta\langle\mathcal{A}\nabla\frac{1}{2}\zeta\rangle + \varepsilon_{\mathcal{F}}. \quad (3.11)$$

In the quantum analogy, \mathcal{F} is momentum and the left hand side of (3.11) is mass times acceleration; the forces are on the right of (3.11). Starting from the end, $\varepsilon_{\mathcal{F}}$ is a dissipative term defined in Appendix B. The third term on the right of (3.11) is the force due to the gradient of the potential $\zeta/2$. The second term on the right of (3.11) represents the combined effect of stretching and tilting of \mathcal{F} by the geostrophic flow. The first term on the right of (3.11) is a ‘‘vortex force,’’ again due to ζ , but perpendicular to \mathcal{F} . The two $\hat{\mathbf{k}}\times$ terms on the right lack quantum analogs.

The results in this section are obtained from the wave equation (2.8) without using the PV equation (2.7). In other words, (3.5)-(3.11) apply to the YBJ equation with e^{imz} structure regardless of the balanced-flow dynamics. We turn now to energy conservation and consideration of the PV equation (2.7).

3.3. Energy conservation and conversion

The energy conservation law is considerably more complicated than action conservation. We sequester the details of the local conservation laws to Appendix B and present here the simpler results obtained by domain averaging those local conservation laws. The results in (3.12) through (3.16) below are obtained by: (1) multiplying the wave equation (2.8) by $\Delta\phi^*$, forming the average $\langle \rangle$, and adding the complex conjugate; and by (2) multiplying the PV equation (2.7) by $-\psi$ and averaging $\langle \rangle$.

For the wave potential energy in (3.4) and the balanced kinetic energy in (3.3) we find

$$\frac{d\langle\mathcal{P}\rangle}{dt} = \Gamma_r + \Gamma_a + \varepsilon_{\mathcal{P}}, \quad (3.12)$$

$$\frac{d\langle\mathcal{K}\rangle}{dt} = -\Gamma_r - \Gamma_a + \Xi + \varepsilon_{\mathcal{K}}, \quad (3.13)$$

where the ‘‘conversion terms’’ in (3.12) and (3.13) are

$$\Gamma_r \stackrel{\text{def}}{=} \left\langle \frac{1}{2}\zeta\nabla\cdot\mathcal{F} \right\rangle, \quad (3.14)$$




FIGURE 3. Illustration of energy conversion terms in the Lamb-Chaplygin dipole solution with parameters presented in table 2. (a) The action flux \mathcal{F} overlain on contours of relative vorticity $\Delta\psi/(U_e k_e) = [-1.5, -0.5, 0.5, 1.5]$, with dashed lines showing negative values; the scale of the action flux is $F = f_0 \lambda^2 k_e U_w^2$. (b) The local advective conversion (i.e., the unaveraged Γ_a) overlain on contours of streamfunction $\psi \times (U_e/k_e) = [-8, -4, -2, 0, 2, 4, 8]$.




FIGURE 4. Diagnostics of the Lamb-Chaplygin dipole solution with parameters presented in table 2. (a) Energy change about initial condition. (b) Wave potential energy budget (3.12).

and

$$\Gamma_a \stackrel{\text{def}}{=} \frac{\lambda^2}{4} \langle \Delta\phi^* J(\psi, \phi) + \Delta\phi J(\psi, \phi^*) \rangle, \quad (3.15)$$

$$= -\frac{\lambda^2}{2} \left\langle \begin{bmatrix} \phi_x^* & \phi_y^* \end{bmatrix} \begin{bmatrix} -\psi_{xy} & \frac{1}{2}(\psi_{xx} - \psi_{yy}) \\ \frac{1}{2}(\psi_{xx} - \psi_{yy}) & \psi_{xy} \end{bmatrix} \begin{bmatrix} \phi_x \\ \phi_y \end{bmatrix} \right\rangle. \quad (3.16)$$

The dissipative terms, $\varepsilon_{\mathcal{P}}$, $\varepsilon_{\mathcal{K}}$ and Ξ are defined in appendix B. Ξ in (3.13) is particularly interesting: dissipation of waves D_ϕ produces balanced kinetic energy. Summing (3.12) and (3.13) the “conversion” terms Γ_r and Γ_a cancel, and we obtain the conservation law for total energy $\langle \mathcal{E} \rangle = \langle \mathcal{P} + \mathcal{K} \rangle$.

The refractive conversion Γ_r stems from $i\phi\zeta/2$ in the wave equation and is easy to interpret: the convergence of the wave action flux, $\nabla \cdot \mathcal{F} < 0$, into anti-cyclones, $\zeta < 0$, is

TABLE 3. The time-integrated budget of wave potential energy and quasigeostrophic kinetic energy of the Lamb-Chaplygin dipole solutions with parameters provided in table 2. The energy budgets close within 10^{-6} %.

$\dot{\mathcal{P}}$ budget	Fractional size ($\int \dot{\mathcal{P}} dt / \Delta \mathcal{P}$)	$\dot{\mathcal{K}}$ budget	Fractional size ($\int \dot{\mathcal{K}} dt / \Delta \mathcal{K}$)
Γ_r	0.228	$-\Gamma_r$	-0.227
Γ_a	0.778	$-\Gamma_a$	-0.774
—	—	Ξ_r	0.004
—	—	Ξ_a	0.0
χ_ϕ	-0.006	ϵ_ψ	-0.003
Res.	0.0	Res.	0.0

a source of wave potential energy \mathcal{P} ; similarly, the divergence of the wave action flux from cyclones is a source of \mathcal{P} . Figure 3a shows the convergence of \mathcal{F} into the anti-cyclone (and divergence from the cyclone) of the dipole solution at $t \times U_e k_e = 1$, which yields the sharp initial increase of $\langle \mathcal{P} \rangle$ discussed below. Ehrenfest's theorem in (3.11) illuminates the initial structure of \mathcal{F} in figure 3. Because ϕ is initially uniform, the initial tendency of \mathcal{F} is

$$\partial_t \mathcal{F}(0) = -\eta \mathcal{A} \nabla \frac{1}{2} \zeta. \quad (3.17)$$

Thus the action flux is initially anti-parallel to the gradient of relative vorticity (see figure 3a), and wave action converges into anti-cyclones (and diverges from cyclones).

The advective conversion Γ_a in (3.12) and (3.13) stems from the term $J(\psi, \phi)$ in the wave equation (2.8) and is a source of $\langle \mathcal{P} \rangle$ due to straining and deformation of the wave field by the geostrophic flow. The symmetric 2×2 matrix in (3.16) is the strain or deformation tensor of the geostrophic flow. Thus, in analogy with passive scalar gradient amplification, straining also enhances gradients of the back-rotated near inertial velocity ϕ , thereby generating wave potential energy $\langle \mathcal{P} \rangle$.

3.4. Energetics of the Lamb-Chaplygin dipole solution

Figure 4 shows the energetics of the Lamb-Chaplygin dipole solution in figure 1. In figure 4a, \mathcal{P} increases at the expense of \mathcal{K} , while \mathcal{A} is conserved. The wave potential energy budget in figure 4b shows that this stimulated generation occurs in two stages. First, refraction of the initially uniform wave field causes a dramatic concentration of waves into the anti-cyclone producing a sharp increase \mathcal{P} through Γ_r . But this rapid initial energy conversion does not last long because the wave feedback deforms the anti-cyclone and dispersion radiates waves away from the dipole (see figure 1). Thus in figure 4b, Γ_r decreases sharply, and eventually reverses sign at $t \times U_e k_e \approx 8$.

The second stage of stimulated generation starts after refraction has created dipole-scale waves. Advection by the balanced flow can then strain the waves, further reducing their lateral scale (figure 4b). The ensuing advective conversion, Γ_a , starts at $t \times U_e k_e \approx 4$. Straining by the balanced flow sustains this advective generation of \mathcal{P} . The waves eventually escape the straining regions through dispersion and the conversion nearly halts at $t \times U_e k_e = 30$. The time-integrated Γ_a accounts for $\approx 78\%$ of the wave potential energy generation; table 3 summarizes the energy budget.

3.5. Summary

The expressions for energy conversion in (3.14) and (3.16) clarify the mechanism of stimulated generation triggered by the initially uniform near-inertial wave in (2.14). First, refraction causes a convergence of wave action into anti-cyclones. Then advection strains the waves, reducing their lateral scale. Both processes amplify the lateral gradients of wave amplitude, thereby generating wave potential energy \mathcal{P} at the expense of balanced kinetic energy \mathcal{K} . Wave action \mathcal{A} is conserved throughout this process.

In the remainder of this paper, we describe and quantify stimulated generation in an idealization of an oceanographic post-storm scenario: the uniform initial near-inertial wave in (2.14) interacts with two-dimensional turbulence.

4. 2D turbulence modified by near-inertial waves

To study the energy exchange between near-inertial waves and geostrophic flow in a turbulent regime, we consider a barotropic flow that emerges from random initial conditions integrated for 20 eddy turnover time units. In other words, we first integrate the initial condition

$$\psi(x, y, t \times U_e k_e = -20) = \sum_{k,l} \psi_{\mathbf{k}} \cos(kx + ly + \chi_{\mathbf{k}}) \quad (4.1)$$

with waveless QG dynamics before introducing the wave in (2.14) at $t \times U_e k_e = 0$. Above, $\chi_{\mathbf{k}}$ is a random phase uniformly distributed on $[0, 2\pi)$, and $\psi_{\mathbf{k}}$ is the streamfunction isotropic spectrum

$$\psi_{\mathbf{k}} = C \times \{|\mathbf{k}| [1 + (|\mathbf{k}|/k_e)^4]\}^{-1/2}, \quad (4.2)$$

with the wavenumber magnitude $|\mathbf{k}|^2 = k^2 + l^2$. The prescribed initial energy $U_e^2/2$ determines the constant C :

$$\sum_{k,l} \underbrace{|\mathbf{k}|^2 \psi_{\mathbf{k}}^2}_{\stackrel{\text{def}}{=} \mathcal{K}_e} = \frac{1}{2} U_e^2. \quad (4.3)$$

The kinetic energy spectrum, \mathcal{K}_e , peaks at the energy-containing scale k_e^{-1} . At scales larger than k_e^{-1} , \mathcal{K}_e has a linear dependence on $|\mathbf{k}|$, whereas \mathcal{K}_e decays as $|\mathbf{k}|^{-3}$ at scales smaller than k_e^{-1} . This red spectrum ensures insignificant loss of energy by small-scale dissipation D_q in (2.7). Over the course of the simulations described below, the centroid wavenumber of the balanced kinetic energy spectrum decreases by 50%; k_e^{-1} is thus a reasonable scale to characterize the size of the balanced flow throughout the evolution.

In the case with no waves, that is $q^w = 0$, the PV equation (2.7) reduces to 2D fluid mechanics and the quasi-inviscid evolution of a random initial condition is the well-studied problem of 2D turbulence. Stirring of vorticity $\Delta\psi$ transfers enstrophy towards small scales; energy flows to large scales. Most of enstrophy is dissipated within few eddy turnover times, whereas kinetic energy is nearly conserved. Vorticity concentrates into localized coherent structures: after 20 eddy turnover time units, the vorticity is well-organized into an ensemble of vortices that form via like-sign vortex merging (e.g., [Fornberg 1977](#); [McWilliams 1984](#)).

4.1. Relevant parameters

The scaling

$$\text{length} \sim k_e^{-1}, \quad \text{time} \sim (U_e k_e)^{-1}, \quad \psi \sim U_e k_e^{-1}, \quad \text{and} \quad \phi \sim U_w, \quad (4.4)$$

TABLE 4. Description of parameters of the 2D turbulence simulations. The initial condition have Rossby number $Ro = U_e k_e / f_0 \approx 0.05$, wave dispersivity $\hbar = f_0 \lambda^2 k_e / U_e \approx 0.5 - 2$, and wave amplitude $\alpha = Ro(U_w / U_e)^2 \approx 0.1$.

Parameter	Description	Value
$2\pi k_e^{-1}$	Energy-containing scale	$L_d/10 \approx 125$ km
U_e	Eddy velocity	5×10^{-2} m s $^{-1}$
U_w	NIW speed	1×10^{-1} m s $^{-1}$
N_0	Buoyancy frequency	5×10^{-3} s $^{-1}$
f_0	Coriolis frequency	10^{-4} s $^{-1}$
$2\pi m^{-1}$	NIW vertical wavelength	280 – 560m
κ_e	PV biharmonic diffusivity	5×10^6 m 4 s $^{-1}$
ν_w	NIW biharmonic viscosity	5×10^6 m 4 s $^{-1}$
N	Number of modes	1024
L	Domain size	$2\pi \times 200$ km

TABLE 5. The time-integrated budget of wave potential energy and QG kinetic energy of the reference 2D turbulence solution with parameters in 4. The energy budgets close within 0.1%.

$\dot{\mathcal{P}}$ budget	Fractional size ($\int \dot{\mathcal{P}} dt / \Delta \mathcal{P}$)	$\dot{\mathcal{K}}$ budget	Fractional size ($\int \dot{\mathcal{K}} dt / \Delta \mathcal{P}$)
Γ_r	0.117	$-\Gamma_r$	-0.108
Γ_a	0.907	$-\Gamma_a$	-0.839
–	–	Ξ_r	0.009
–	–	Ξ_a	0.003
χ_ϕ	-0.026	ϵ_ψ	-0.062
Res.	0.003	Res.	0.003

shows that there are two important dimensionless control parameters. The first is

$$\alpha \stackrel{\text{def}}{=} \underbrace{\frac{U_e k_e}{f_0}}_{\stackrel{\text{def}}{=} Ro} \times \left(\frac{U_w}{U_e} \right)^2, \quad (4.5)$$

which scales the contribution of the wave terms in the potential vorticity (2.6). The second dimensionless parameter is

$$\hbar \stackrel{\text{def}}{=} \eta \times \frac{k_e}{U_e} = Ro^{-1} \times (\lambda k_e)^2, \quad (4.6)$$

which scales wave dispersion against the effects of advection and refraction. Assuming that the wave horizontal scale is k_e^{-1} , $(\lambda k_e)^2$ is the wave Burger number, which is small for near-inertial waves.

4.2. Solution with $\hbar = 1$ and $\alpha = 0.1$

Figure 5 shows snapshots of a solution with $\hbar = 1$ and $\alpha = 0.1$ and further parameters in table 4. This turbulence solution shares qualitative aspects of the Lamb-Chaplygin solution. Starting from a uniform wave field in (2.14), refraction quickly concentrates the waves into anti-cyclones. Initially the action density \mathcal{A} is uniform but by $t \times U_e k_e \approx 1$,




FIGURE 5. Snapshots of the turbulence solution with parameters in table 4. Upper panels: PV q . Middle panels: wave kinetic energy density $|\phi|^2$. Bottom panels: wave buoyancy, with scale $B = k_e m U_w f_0 \lambda^2$. These plots show $(1/2)^2$ of the domain.

\mathcal{A} varies on eddy scales by a factor of two with significant focussing of waves, indicated by maxima of \mathcal{A} , into anti-cyclones (compare middle and upper panels of figure 5).

Dispersion radiates waves from the vortices; advection enhances the gradients of back-rotated velocity ϕ (see lower panels of figure 5, which depict wave buoyancy). By $t \times U_e k_e \approx 10$, \mathcal{A} varies by a factor of five and the wave buoyancy is amplified by a factor of two. The evolution of potential vorticity q is similar to that in the waveless problem: like-sign vortices merge into bigger vortices. The big vortices keep straining the waves, generating smaller scales in the wave field.

Figure 6a shows the inexorable increase in wave potential energy $\langle \mathcal{P} \rangle$ and the corresponding decrease in balanced kinetic energy $\langle \mathcal{K} \rangle$. In figure 6b quick wave refraction results in an initial sharp generation of $\langle \mathcal{P} \rangle$ at the expense of balanced kinetic energy $\langle \mathcal{K} \rangle$. As in the Lamb-Chaplygin solution, the positive refractive conversion, $\Gamma_r > 0$, is ephemeral: in figure 6b, Γ_r peaks at $t \times U_e k_e \approx 2$ and then decays rapidly, eventually changing sign at $t \times U_e k_e \approx 5$. But a significant positive advective conversion, $\Gamma_a > 0$, sustains stimulated generation so that $\langle \mathcal{P} \rangle$ ultimately increases approximately linearly with time.




FIGURE 6. Diagnostics of the 2D turbulence solution with parameters presented in table 4. (a) Energy change about initial condition. (b) Wave potential energy budget (3.12).

After 25 eddy-turnover time units, the balanced kinetic energy $\langle \mathcal{K} \rangle$ has decayed by about 14% from its initial value. Most of this loss is by stimulated generation of $\langle \mathcal{P} \rangle$. As in the Lamb–Chaplygin solution, advective conversion Γ_a accounts for most of the energy change. Table 5 presents further details of the energy budget.

The solution illustrates interesting characteristics of stimulated generation. First, the role of refraction is catalytic in that it generates the initial eddy-scale gradients in ϕ that are then enhanced by advective straining; the advective conversion, Γ_a in (3.16), ultimately accounts for most of the energy transfer from turbulence to waves. Second, the roughly linear-in-time growth of wave potential energy $\langle \mathcal{P} \rangle$ is very slow in comparison with exponential increase of passive-scalar tracer gradients in turbulent velocity fields. The relatively slow growth of $\langle \mathcal{P} \rangle$ suggests that wave dispersion plays an important role in slowing and perhaps opposing advective straining (see section 5 for further discussion of dispersion and “wave escape”). To investigate whether these characteristics are general we consider solutions with varying vertical wavelengths and therefore different dispersivities.

4.3. Varying dispersivity

Figure 7 shows snapshots of potential vorticity q and its constituents in a set of solutions with varying the vertical wavelength $2\pi m^{-1}$ from 280 to 560 m, yielding dispersivities ranging from 0.5 to 2. (All other parameters are fixed.) The potential vorticity q shows more small-scale filamentation with decreasing dispersivity, but it is otherwise similar across the three solutions. The partition into relative vorticity $\Delta\psi$ and wave potential vorticity q^w , however, depends significantly on dispersivity. In particular, q^w develops smaller scales and larger amplitudes with decreasing dispersivity. As anticipated by the dipole example in figure 2, there is cancellation of small-scale features in q^w against those in ζ so that q is relatively smooth even in the solution with weak dispersion $\bar{h} = 0.5$.

The initial evolution of the uniform wave field is similar across dispersivities, with refraction initially generating eddy-scale gradients of the waves—see figure 8. Refraction produces a sharp initial increase of wave potential energy and decrease of balanced kinetic energy, which is almost independent of dispersivity. Figure 8c shows that this initial




FIGURE 7. Snapshots of PV q and its decomposition into relative vorticity $\zeta = \Delta\psi$, and wave potential vorticity q^w . The snapshots were taken at $t \times U_e k_e = 25$.

“refractive stage” yields a strongly negative wave-vorticity correlation r ,

$$r \stackrel{\text{def}}{=} \frac{\langle \zeta \mathcal{A}' \rangle}{\sqrt{\langle \zeta^2 \rangle \langle \mathcal{A}'^2 \rangle}}, \quad (4.7)$$

where $\mathcal{A}' \stackrel{\text{def}}{=} (|\phi|^2 - \langle |\phi|^2 \rangle) / f_0$; in figure 8c the early negative r is nearly independent of dispersivity. Because significant energy exchange takes place in the anti-cyclones due to the initial wave concentration, a positive vorticity skewness ensues (figure 8d). Once the eddy-scales are created, advection strains the waves and generates further wave potential energy at the expense of balanced kinetic energy. It is in this stage that the dependence on dispersivity is pronounced: weakly dispersive waves are strained further than strongly dispersive waves. Thus the advective conversion becomes stronger with decreasing dispersivity (figure 8b). Advection and dispersion significantly reduces the wave-vorticity correlation; the reduction in correlation increases as the dispersivity decreases (figure 8c). For the weakest dispersivity considered, the wave-vorticity correlation becomes weakly positive likely because of the early positive vorticity skewness.




FIGURE 8. The energetics of 2D turbulence solutions with different dispersivities. (a) Energy change about the initial condition. (b) The energy conversion terms in (3.12). (c) The correlation between relative vorticity and wave kinetic energy. (d) The skewness of relative vorticity.




FIGURE 9. Energy-preserving spectra of 2D turbulence solutions with different dispersivities. The three panels show spectra of balanced kinetic energy \mathcal{K} , wave action \mathcal{A} , and wave potential energy \mathcal{P} . All solid lines correspond to spectra at $t \times U_e k_e = 25$ and the dashed line in \mathcal{K} is the balanced kinetic energy spectrum at $t \times U_e k_e = 0$. All spectra are normalized by their total energy, i.e., the area under each curve is one.

In all solutions reported above, the evolution of the balanced flow is similar to that of waveless 2D turbulence: there is a transfer of balanced energy towards larger scales driven by merger of like-signed vortices—see the left panel of figure 9. The main difference is that balanced kinetic energy is constantly transformed into wave potential energy via stimulated generation. The stimulated generation process is associated with a forward transfer of wave action \mathcal{A} from the infinite horizontal scale in the initial condition (2.14) to the eddy scale; see the middle panel of figure 9. The wave potential energy density \mathcal{P} on the right panel of figure 9 develops significantly smaller scales than those of the balanced kinetic energy \mathcal{K} .

5. Wave escape

The expression for the energy conversion in (3.12) illuminates the physics of stimulated generation: both convergence of wave action density into anti-cyclones (3.14) and geostrophic straining of the wave field (3.16) are sources of wave potential energy and sinks of balanced kinetic energy. But this characterization of stimulated generation ignores the important role of wave dispersion—waves can propagate out of the vorticity or straining regions, thereby reducing the correlations Γ_r and Γ_a required for stimulated generation.

Wave dispersion is the only mechanism that upsets stimulated generation in the quasi-inviscid solutions described in this paper. In all solutions, after an initial conversion due to refraction, advective straining accounts for most of the energy conversion. Experience with the passive-scalar problem suggests (incorrectly) that the wave potential energy \mathcal{P} should then increase exponentially with time as $\nabla\phi$ is amplified by stirring (e.g., Young *et al.* 1982). But even in the weakly dispersive limit, the waves do not behave as a passive scalar and stimulated generation is much less effective than suggested by this “passive-scalar thinking.” This is because advective straining can only increase $\nabla\phi$ so much: the near-inertial generalized group velocity is $\eta\nabla\Theta$ (cf. section 3), where Θ is the phase of the near-inertial back-rotated velocity $\phi = |\phi|e^{i\Theta}$. Geostrophic straining enhances $\nabla\Theta$ thereby increasing the near-inertial group velocity so that the waves escape the straining region. Thus, straining by a barotropic balanced flow results in near-inertial “wave escape,” as opposed to the “wave capture” described by Bühler & McIntyre (2005). Indeed, wave capture requires *both* lateral strain *and* vertical shear: the vertical-plane-wave model has no vertical shear and therefore wave capture is inoperative; see Thomas (2012) for further discussion of the importance of vertical shear to wave capture.

5.1. Strain flow

We are surprised by the successful resistance mounted by the waves to strain-driven exponential amplification of $\nabla\phi$ and thus seek to illustrate wave escape with simple flows. We first consider the straining flow $\psi = -\alpha xy$. Ignoring dissipation for simplicity, the wave equation (2.8) reduces to

$$\phi_t + \alpha x \phi_x - \alpha y \phi_y - \frac{i}{2} \eta \Delta \phi = 0, \quad (5.1)$$

and the action equation is

$$\mathcal{A}_t + \alpha(x\mathcal{A})_x - \alpha(y\mathcal{A})_y + \nabla \cdot \mathcal{F} = 0. \quad (5.2)$$

Without vorticity and dissipation, Ehrenfest’s theorem (3.11) reduces to

$$\partial_t [\langle \mathcal{F}^x \rangle, \langle \mathcal{F}^y \rangle] = [-\alpha \langle \mathcal{F}^x \rangle, +\alpha \langle \mathcal{F}^y \rangle], \quad (5.3)$$




FIGURE 10. The escape of a Gaussian near-inertial wave packet from the saddle of a strain flow. The Gaussian decay scale is $\mu = \nu$. The waves are weakly dispersive: $\hbar = \eta/\alpha\mu^2 \approx 0.1$. Black contours show the streamfunction $\psi = -\alpha xy$, and colors represent wave zonal velocity; the colorbar limits are fixed. The black dot indicates the center of the packet and the gray line tracks its trail.

with solution

$$[\langle \mathcal{F}^x \rangle, \langle \mathcal{F}^y \rangle] = [\langle \mathcal{F}^x \rangle_0 e^{-\alpha t}, \langle \mathcal{F}^y \rangle_0 e^{\alpha t}], \quad (5.4)$$

where the subscript 0 denotes the initial condition. For a compact wave packet, with a well-defined uniform group velocity \mathbf{c}_g , the action flux is

$$\langle \mathcal{F} \rangle = \mathbf{c}_g \langle \mathcal{A} \rangle. \quad (5.5)$$

Because $\langle \mathcal{A} \rangle$ is constant, (5.4) is thus also a solution for $\mathbf{c}_g(t)$.

The position of the center of the packet is

$$\mathbf{x}_c(t) \stackrel{\text{def}}{=} \langle \mathbf{x} \mathcal{A} \rangle / \langle \mathcal{A} \rangle. \quad (5.6)$$

Multiplying (5.2) by \mathbf{x} , averaging, and using (5.4), yields

$$\partial_t x_c - \alpha x_c = c_{g0}^x e^{-\alpha t}, \quad \text{and} \quad \partial_t y_c + \alpha y_c = c_{g0}^y e^{\alpha t}, \quad (5.7)$$

with solution

$$\mathbf{x}_c(t) = \mathbf{x}_{c0} + \mathbf{c}_{g0} \frac{\sinh \alpha t}{\alpha}. \quad (5.8)$$

The trajectory of the packet is therefore a straight line following the initial group velocity \mathbf{c}_{g0} . The wave packet is not deflected by the hyperbolic streamlines of the straining flow and the packet escapes by accelerating exponentially with time along a straight-line trajectory. This requires that the group velocity $\mathbf{c}_g(t)$ adjusts in magnitude and direction so as to keep the center of the packet on the straight and narrow.

The result in (5.8) is so remarkable that it is reassuring to obtain it without invoking Ehrenfest's theorem. We thus consider the specific example of a wave packet launched in the strain field with the Gaussian initial condition

$$\phi(x, y, 0) = \frac{2\pi}{\mu\nu} \exp \left[-\frac{1}{2} \left(\frac{x^2}{\mu^2} + \frac{y^2}{\nu^2} \right) + i(px + qy) \right]. \quad (5.9)$$

If $p\mu \gg 1$ and $q\nu \gg 1$ this is a wave packet with initial wavenumber (p, q) and initial group velocity

$$\mathbf{c}_{g0} = \eta(p, q). \quad (5.10)$$




FIGURE 11. A comparison of passive-scalar and wave solutions ($\hbar \approx 0.09$) with same initial conditions (the wave kinetic energy is equal to the passive-scalar variance) and the same small-scale dissipation. (a) Initial condition of wave back-rotated zonal velocity and of the passive scalar. (b) Wave back-rotated zonal velocity at $t \times U_e k_e = 10$. (c) Passive-scalar concentration at $t \times U_e k_e = 10$. (d) Variance of wave velocity or passive-scalar variance. (e) Variance of wave velocity gradient or passive scalar. In (d) and (e), the diagnostics are normalized by their initial values.

The exact solution to (5.1), subject to the initial condition (5.9), is

$$\phi = \frac{2\pi}{\sqrt{(\mu^2 + i\eta f)(\nu^2 + i\eta g)}} \times \exp \left[-\frac{1}{2} \frac{x^2 e^{-2\alpha t} - 2i\mu^2 p x e^{-\alpha t} + i\eta f \mu^2 p^2}{\mu^2 + i\eta f} - \frac{1}{2} \frac{y^2 e^{2\alpha t} - 2i\nu^2 q y e^{\alpha t} + i\eta g \nu^2 q^2}{\nu^2 + i\eta g} \right], \quad (5.11)$$

with

$$f \stackrel{\text{def}}{=} \frac{1 - e^{-2\alpha t}}{2\alpha} \quad \text{and} \quad g \stackrel{\text{def}}{=} \frac{e^{2\alpha t} - 1}{2\alpha}. \quad (5.12)$$

Figure 10 illustrates the solution (5.11). The strain flow tilts the packet to align it with the x -axis. The flow then exponentially stretches the packet, which in turn escapes from the saddle point. To calculate the trajectory of the packet we note that

$$|\phi|^2 = \frac{(2\pi)^2}{\sqrt{(\mu^4 + \eta^2 f^2)(\nu^4 + \eta^2 g^2)}} \exp \left[-\frac{\mu^2 (x e^{-\alpha t} - \eta f p)^2}{\mu^4 + \eta^2 f^2} - \frac{\nu^2 (y e^{\alpha t} - \eta g q)^2}{\nu^4 + \eta^2 g^2} \right], \quad (5.13)$$

and we recover $\mathbf{x}_c(t)$ in (5.8) as the center of the packet in (5.13).

5.2. A flow with strain and vorticity

As a second example of wave escape, figure 11 shows a numerical solution for a wave packet launched at the saddle point of a large-scale balanced flow with $\psi = \sin x + \sin y$. The small-time evolution of the packet is predicted by the strain-flow solution discussed above. The top row of figure 11 shows that the behavior of the wave packet is qualitatively different from that of a passive scalar in the same flow: the passive scalar is stretched out along the separatrices while the wave packet escapes into the vortex centers. Thus the passive scalar packet is strained and quickly diffused into oblivion. On the other hand, the waves are strained just so much, resulting in acceleration and escape from the straining region; the waves finally concentrate in the regions with non-zero vorticity, i.e., in the regions where the Okubo-Weiss criterion indicates no exponential stretching.

On the bottom row, figure 11d shows that while wave action $\propto |\phi|^2$ is nearly conserved, the analogous passive-scalar variance is strongly dissipated. Figure 11e shows that the variance of the passive-scalar gradient at first increases exponentially due to straining and then decays due to diffusion. On the other hand, the potential energy of the waves $\propto |\nabla\phi|^2$ increases slowly and then oscillates around an equilibrium level. The wave-escape phenomenology in the turbulence solutions of section 4 qualitatively resembles that seen in this simple flow. In particular, the wave potential energy does not reach the dissipative scale (figure 9c).

6. The Eulerian-mean viewpoint

In section 3 we developed conservation laws for \mathcal{A} , \mathcal{P} and \mathcal{K} based on the Lagrangian-mean streamfunction ψ and the backrotated velocity ϕ . Material conservation of PV is central to the NIW-QG system and thus the *Lagrangian-mean velocity*, $(-\psi_y, \psi_x)$, must be a key field. But then $\mathcal{K} = |\nabla\psi|^2/2$ is not the kinetic energy of the *Eulerian-mean velocity* and therefore Reynolds stress diagnostics do not directly indicate changes in \mathcal{K} . In fact Reynolds stresses have not been mentioned in section 3 and one might wonder how these standard statistics are related to the NIW-QG energy conversions Γ_r and Γ_a ? In this section we identify the Eulerian-mean energy transfers in terms of NIW-QG fields.

6.1. Kinetic energy of the Lagrangian-mean and Eulerian-mean flows

Appendix A.2 shows that in the vertical-plane-wave model, the Stokes drift is horizontally non-divergent, with streamfunction $-\mathcal{A}$. Thus the Stokes velocity is

$$\mathbf{u}^S = -\hat{\mathbf{k}} \times \nabla\mathcal{A}, \quad (6.1)$$

and the Eulerian-mean streamfunction is

$$\psi^E \stackrel{\text{def}}{=} \psi + \mathcal{A}. \quad (6.2)$$

To illustrate the important differences between Lagrangian-mean and Eulerian-mean viewpoints, figure 12 shows a snapshot of the dipole example from section 2.3. This snapshot was taken at $t \times U_e k_e = 10$, just after the end of the refractive stage of energy conversion, when waves are strongly concentrated in the anti-cyclone. While ψ is fairly symmetric, the Eulerian-mean streamfunction ψ^E displays a stronger anti-cyclone. The Stokes drift is concentrated in the negative ζ region and is anti-parallel to the Eulerian-mean flow. Thus the asymmetry in ψ^E is compensated by a strong ‘‘Stokes cyclone,’’ which is set up during the refractive stage and thus the Lagrangian-mean streamfunction ψ is more nearly a symmetric dipole.




FIGURE 12. Snapshot of the dipole Lagrangian-mean streamfunction and its decomposition into Eulerian-mean and Stokes drift. The snapshots are at $t \times U_e k_e = 10$, just after the end of the refractive stage. The spatial average of each field was removed for direct comparison.

Using the decomposition (6.2), the balanced kinetic energy $\mathcal{K} = |\nabla\psi|^2/2$ is

$$\mathcal{K} = \underbrace{\frac{1}{2}|\nabla\psi^E|^2}_{\stackrel{\text{def}}{=} \mathcal{K}^E} + \underbrace{\frac{1}{2}|\nabla\mathcal{A}|^2 - \nabla\psi^E \cdot \nabla\mathcal{A}}_{\stackrel{\text{def}}{=} \mathcal{K}^S}. \quad (6.3)$$

\mathcal{K} does not diagonalize, i.e., the cross-term $\nabla\psi^E \cdot \nabla\mathcal{A}$ is non-zero, and the “Stokes kinetic energy” \mathcal{K}^S is not sign definite.

The first goal here is to obtain an expressions for the time rate of change of \mathcal{K}^S and \mathcal{K}^E . We begin with yet another \mathcal{A} - \mathcal{F} identity: using the definitions of \mathcal{A} and \mathcal{F} in (3.1) and (3.6) yields

$$\mathcal{A} \nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) = -\mathcal{F} \cdot \hat{\mathbf{k}} \times \nabla \mathcal{A}. \quad (6.4)$$

The average of (6.4), combined with a standard vector identities, results in

$$\langle \mathcal{A} \nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) \rangle = \langle \mathcal{F} \cdot \hat{\mathbf{k}} \times \nabla \mathcal{A} \rangle = 0. \quad (6.5)$$

Equation (6.5) implies that the Stokes velocity \mathbf{u}^S is, on average, orthogonal to the action flux \mathcal{F} .

Forming $\langle \mathcal{A}q \rangle$, and combining the expression for q^w in (3.9) with the identity (6.5), yields

$$\langle \mathcal{K}^S \rangle = \langle \mathcal{A}q \rangle. \quad (6.6)$$

An expression for the rate of change of $\langle \mathcal{K}^S \rangle$ follows by combining the PV advection equation (2.7) with the action conservation (3.5):

$$\frac{d\langle \mathcal{K}^S \rangle}{dt} = -\Gamma_S + \underbrace{\langle \mathcal{A}Dq \rangle + \langle qD\mathcal{A} \rangle}_{\stackrel{\text{def}}{=} \varepsilon_S}, \quad (6.7)$$

where

$$\Gamma_S \stackrel{\text{def}}{=} \langle q \nabla \cdot \mathcal{F} \rangle. \quad (6.8)$$

Finally we obtain the rate of change of the Eulerian-mean kinetic energy by combining (6.7) with (3.13):

$$\frac{d\langle \mathcal{K}^E \rangle}{dt} = \Gamma_S - \Gamma_r - \Gamma_a + \Xi + \varepsilon_{\mathcal{K}} - \varepsilon_S. \quad (6.9)$$




FIGURE 13. Diagnostics of the 2D turbulence solution with parameters presented in table 4. (a) Kinetic energy change about initial condition. (b) Refractive and advective conversion terms, and shear production.

6.2. Reynolds stresses and buoyancy fluxes

Within the Eulerian framework the horizontal velocity is represented as

$$\mathbf{u} = \mathbf{u}^E + \tilde{\mathbf{u}}, \quad (6.10)$$

where superscript E denotes the Eulerian-mean and tilde denotes the near-inertial wave velocity. With this decomposition, the Eulerian-mean velocity satisfies

$$\mathbf{u}_t^E + \mathbf{u}^E \cdot \nabla \mathbf{u}^E + (\tilde{\mathbf{u}} \cdot \nabla \tilde{\mathbf{u}})^E + \hat{\mathbf{k}} \times f_0 \mathbf{u}^E + \nabla p^E = 0. \quad (6.11)$$

(Dissipation is neglected in the following discussion.) As in the vertical plane-wave model, we confine attention to mean velocities independent of z , so that $\mathbf{u}^E = (-\psi_y^E, \psi_x^E)$ where ψ^E is the Eulerian-mean streamfunction introduced in (6.2). Forming $\langle \mathbf{u}^E \cdot (6.11) \rangle$ yields

$$\frac{d\langle \mathcal{K}^E \rangle}{dt} = \underbrace{\langle (\tilde{v}^2 - \tilde{u}^2)^E \psi_{xy}^E + (\tilde{u}\tilde{v})^E (\psi_{xx}^E - \psi_{yy}^E) \rangle}_{\stackrel{\text{def}}{=} RSP}, \quad (6.12)$$

Above RSP is the Reynolds shear production of Eulerian-mean kinetic energy. Comparing (6.12) with (6.9), and ignoring the dissipative contributions, we see that

$$RSP = \Gamma_S - \Gamma_r - \Gamma_a. \quad (6.13)$$

We can also consider the source of near-inertial potential energy by starting with the wave buoyancy equation

$$\tilde{b}_t + \mathbf{u}^E \cdot \nabla \tilde{b} + \nabla \cdot [\tilde{\mathbf{u}} \tilde{b} - (\tilde{\mathbf{u}} \tilde{b})^E] + \tilde{w} N^2 = 0, \quad (6.14)$$

where we have confined attention to the vertical-plane-wave model so that $b^E = 0$. Multiplying (6.14) by \tilde{b} , taking the Eulerian average, and then the domain average $\langle \rangle$,

we have

$$\frac{d\langle\mathcal{P}\rangle}{dt} = - \underbrace{\langle(\tilde{w}\tilde{b})^E\rangle}_{\stackrel{\text{def}}{=}BF}, \quad (6.15)$$

$$= \Gamma_r + \Gamma_a. \quad (6.16)$$

Above $\mathcal{P} = (\tilde{b}^2)^E/2N^2$ is the wave potential energy and BF is the buoyancy flux; we have used (3.12) to relate Γ_r and Γ_a to the Eulerian-mean buoyancy flux.

Finally, from balanced energy equation (3.13) and $\mathcal{K} = \mathcal{K}^E + \mathcal{K}^S$, we deduce that

$$\frac{d\langle\mathcal{K}^S\rangle}{dt} = BF - RSP, \quad (6.17)$$

$$= -\Gamma_S. \quad (6.18)$$

Figure 13a shows the decomposition of $\langle\mathcal{K}\rangle$ into $\langle\mathcal{K}^E\rangle$ and $\langle\mathcal{K}^S\rangle$ for the main 2D turbulence solution discussed in section 4. In the initial condition there is no Stokes flow and $\mathcal{K} = \mathcal{K}^E$. Then in the refractive stage, the kinetic energy of the Eulerian-mean flow $\langle\mathcal{K}^E\rangle$ increases, while kinetic energy of the Lagrangian-mean flow $\langle\mathcal{K}\rangle$ decreases; the ‘‘Stokes energy’’, $\mathcal{K}^S = \mathcal{K} - \mathcal{K}^E$, is initially zero and becomes negative, due to a positive ψ^E - \mathcal{A} correlation (cf. figure 12). As illustrated in figure 12, the refractive stage, which creates the strong spatial modulations in action density \mathcal{A} , can also be viewed as the setting-up the Stokes velocity in (6.1).

In the refractive stage, the advective conversion Γ_a is small and $\Gamma_S \approx 2\Gamma_r$. Hence the shear production is due to the refractive conversion, $RSP \approx \Gamma_r$ (see figure 13b). At later times, during the advective stage, $\Gamma_S - \Gamma_r$ is small, and therefore $RSP \approx BF \approx -\Gamma_a$. At this stage, $\partial_t\langle\mathcal{K}\rangle \approx \partial_t\langle\mathcal{K}^E\rangle$, and the connection between the Eulerian-mean and Lagrangian-mean viewpoints is straightforward: the kinetic energy extracted from the Eulerian-mean flow via Reynolds shear production approximately matches the creation of wave potential energy through buoyancy fluxes. In general, however, the connection is convoluted, involving the rate of change of the ‘‘Stokes energy’’ in (6.17).

7. Discussion and conclusions

7.1. Absence of a direct cascade of wave energy

The solutions reported here introduce the waves at $t = 0$ in (2.14) with infinite spatial scale. Wave refraction, $i\phi\zeta/2$ in (2.8), immediately transfers wave energy to the smaller scales of the balanced relative vorticity ζ . This giant leap across wavenumbers is not a direct cascade of wave energy in the sense of Kolmogorov. And because of wave escape, the wave energy that is so efficiently transferred to eddy scales by refraction does not undergo a turbulence-driven direct cascade to the small length scales at which the wave dissipation in (2.9) is effective. This conclusion hinges on the assumption of a barotropic balanced flow and fixed vertical wavenumber. Similar to the vertical-plane-wave model, shallow-water models lack wave capture and an efficient direct cascade of wave energy (e.g., McIntyre 2009). In less idealized models with balanced baroclinic shear, the vertical wavenumber can increase, and a direct cascade, perhaps resulting in wave capture, can ensue; stimulated generation might then be much stronger than in the vertical-plane-wave model considered here.

7.2. Regimes of wave-modified turbulence

Geostrophic straining accounts for most of the stimulated generation of wave energy in the examples considered in this paper. But refraction plays a fundamental role in these solutions with the uniform initial wave velocity in (2.14), because refraction creates the initial gradients of wave velocity that are then enhanced by geostrophic straining. We experimented by changing the initial condition of ϕ to an eddy-scale plane wave and repeated all the 2D turbulence solutions; the different initial condition significantly suppresses the initial refraction stage, but otherwise yields long-term solutions that are qualitatively similar to the solutions discussed above. Thus, to the extent that the uniform-wave initial condition (2.14) idealizes the generation of large-scale upper-ocean inertial oscillations by storms (e.g., Moehlis & Llewellyn Smith 2001; Danioux *et al.* 2015), the initial refraction is a loss of lateral coherence, or a type of inertial pumping (Young & Ben Jelloul 1997; Klein *et al.* 2004), which is accompanied by an extraction of energy from the balanced flow by the waves.

Although 10-20% of the balanced kinetic energy is converted into wave potential energy, and despite the wave breakage of the symmetry between cyclones and anti-cyclones, the wave-modified turbulence in section 4 remarkably resembles waveless two-dimensional turbulence (e.g., McWilliams 1984): we still observe robust vortices and an increase in vortex length scale due to merger of like-signed vortices. Figure 7 shows small changes in the potential vorticity q and much larger changes in the wave PV q^w induced by changing the dispersivity. In this sense, the turbulent evolution is insensitive to wave modification.

To see significant wave modification of the turbulence we increased the amplitude of the initial wave in (2.14) so that $U_w = 6U_e$ (in section 4, $U_w = 2U_e$). With this level of wave energy the wave-modified 2D turbulence differs qualitatively from the waveless variety (not shown). The potential vorticity develops highly filamentary structures with little vortex formation; this inhibition of vortex formation is stronger in the weakly dispersive limit.

7.3. Energy transfers can be bi-directional in non-turbulent balanced flows

In all solutions considered in sections 2 and 4, the energy transfer is always from the balanced flow into the waves. This positive energy conversion, $\Gamma_r + \Gamma_a > 0$, remains true for very long turbulence simulations (not shown), because the refractive conversion is small at large time and turbulent stirring always increases lateral wave gradients: $\Gamma_r + \Gamma_a \approx \Gamma_a > 0$.

But the energy transfers can be bi-directional for non-turbulent flows. To illustrate this process, we consider a solution with initially uniform wave and $\psi(\mathbf{x}, 0) = \sin x + \sin y$. This non-turbulent balanced flow consists of two vortices of opposite signs; there can be no inverse cascade from this initial condition because $\psi(\mathbf{x}, 0)$ is already at the domain scale. Starting from the initial uniform wave, the refractive conversion generates wave potential energy \mathcal{P} at the expense of balanced kinetic energy \mathcal{K} (figure 14). After this initial refractive stage, however, \mathcal{P} and \mathcal{K} oscillate about equilibrium levels and there are quasi-periodic energy transfers back and forth between \mathcal{P} and \mathcal{K} . In this solution, the refractive conversion Γ_r accounts for most of the quasi-periodic exchanges between \mathcal{K} and \mathcal{P} .

7.4. The correlation of wave amplitude with relative vorticity

A secondary result is the strong time-dependence of the correlation r , defined in (4.7), between incoherent waves and the relative vorticity; r measures the concentration of waves into cyclones or anti-cyclones (Danioux *et al.* 2015). Refraction concentrates




FIGURE 14. Diagnostics of an illustrative solution with initially uniform wave and $\psi(t=0) = \sin x + \sin y$, with $\alpha = 0.1$ and $\bar{h} = 2$. (a) Energy difference about initial condition. (b) Wave potential energy budget.

waves into anti-cyclones and expels them from cyclones, thereby generating an initial strong negative r (see figure 8c). And Ehrenfest’s theorem provides perhaps the simplest explanation for this concentration—see (3.17). As conjectured by [Danioux *et al.* \(2015\)](#), the subsequent return of r towards zero (and even to positive values in the case $\bar{h} = 0.5$) is partially due to the unsteady geostrophic advection. The NIW-QG coupling compounds the unsteady advection: the dramatic initial concentration of waves into anti-cyclones weakens those vortices, with ensuing development of positive skewness of relative vorticity (see figure 8d); the vorticity skewness increases with decreasing dispersivity because weakly dispersive waves extract more energy from the balanced flow (see figure 8).

7.5. Reconciling action conservation with RSP of wave kinetic energy

A main difficulty in making a connection between the NIW-QG model and earlier studies, such as those in table 1, is that the theory uses the Lagrangian-mean geostrophic flow as a primary variable. Previous studies generally employ an Eulerian wave-mean decomposition. [Taylor & Straub \(2016\)](#), for example, use frequency filtering to separate low frequency motions (Eulerian-mean) from high-frequency flow (dominated by near-inertial waves). Taylor & Straub find energy exchange between low-frequency (eddy) kinetic energy and high frequency (near-inertial) kinetic energy resulting from both vertical and horizontal Reynolds stresses; Taylor & Straub refer to this energy transfer as the “advective sink”, meaning a sink of low-frequency, Eulerian-mean kinetic energy. The Reynolds shear production RSP in (6.12) is analogous to the horizontal part of the advective sink.

The Reynolds stresses diagnostics of prior studies in table 1 might seem incompatible with fundamental aspects of the NIW-QG system. For example, the wave action \mathcal{A} in (3.1) differs from the wave kinetic energy $|\phi|^2/2$ only by the constant factor f_0 . Thus NIW-QG action conservation seems inconsistent with Reynolds stress transfer of kinetic

energy from eddies to waves. But this impression is incorrect: Reynolds stresses aren't inconsistent with conservation of \mathcal{A} , nor with stimulated generation.

To understand this, note that conservation of \mathcal{A} in (3.10) is a statement about the *leading order* wave velocities encoded in ϕ ; the balanced kinetic energy \mathcal{K} also involves *higher order* wave kinetic energy in the form of the ‘‘Stokes kinetic energy’’ \mathcal{K}^S : see (6.3) and the surrounding discussion. A main point from section 6.2 is that the energy transferred from the Eulerian-mean velocity by *RSP* is accounted for by \mathcal{K}^S (rather than the leading-order wave kinetic energy $f_0\mathcal{A}$).

At first glance it might seem that because stimulated generation results in an increase in wave potential energy \mathcal{P} , it must necessarily involve a transfer from eddy potential energy. The vertical-plane-wave model is a counterexample: the eddies are barotropic and there is therefore no eddy potential energy to transfer. Indeed, if the eddies are barotropic, then *RSP* is the *only* pathway between the Eulerian-mean kinetic energy \mathcal{K}^E and wave energy. This rather obvious fact is deeply hidden by the Lagrangian-mean average: the *RSP* transfer out of \mathcal{K}^E passes first through \mathcal{K}^S on the way to \mathcal{P} . The intermediate passage through \mathcal{K}^S is an inevitable aspect of the Lagrangian-mean formulation of the NIW-QG system. All of these transfers can be diagnosed using Γ_a , Γ_r and Γ_S and (6.13) relates Reynolds shear production to the three Γ 's.

A full investigation of NIW-QG energetics, including diagnosis of Γ_r , Γ_a and Γ_S in solutions of the Boussinesq equations, is beyond the scope of this article. But we emphasize a crucial simplifying feature of the NIW-QG approximation: there are no important Stokes corrections to pressure and buoyancy, and therefore, to leading order, the Lagrangian-mean velocity is equal to the balanced velocity based on an Eulerian-mean pressure field. In principle, this is a straightforward way of diagnosing Lagrangian-mean velocities from numerical solutions of the Boussinesq equations.

7.6. Final remarks

There are many caveats to the application of our results to the post-storm oceanographic problem. Notably, the lack of geostrophic vertical shear suppresses important mechanisms of vertical refraction and straining, which introduce interesting modifications of the near-inertial wave physics (e.g., Thomas 2012) and can produce strong energy extraction by near-inertial waves (Shakespeare & Hogg 2017). And our focus on quasi-inviscid initial value problems downplays the role of dissipation, but in forced-dissipative solutions, wave dissipation likely controls the strength of stimulated generation. Finally, better understanding the connection between numerical modeling studies (e.g. Taylor & Straub 2016) and the NIW-QG theory (XV; Wagner & Young 2016) deserves further investigation. We hope to explore these topics in future work.

We thank three anonymous referees for constructive criticism and suggestions. This study was supported by the National Aeronautics and Space Administration (NNX16AO5OH) and the National Science Foundation (OCE1357047).

Appendix A. Details of the NIW-QG model

A.1. The NIW-QG model

Using multiscale asymptotic theory, Wagner & Young (2016) derive a model for the coupled evolution of QG balanced flow, near-inertial waves and their second harmonic. Assuming that the second harmonic is zero ($B = 0$), the Wagner & Young (2016) coupled model recovers the XV model in the limit where the waves have vertical scales much

smaller than the balanced flow. In [Wagner & Young \(2016\)](#), the PV is

$$q = (\Delta + L)\psi + \beta y + \frac{1}{2f_0} \left[\Delta \frac{1}{2} |LA|^2 + iJ(LA^*, LA) \right], \quad (\text{A } 1)$$

where $\Delta \stackrel{\text{def}}{=} \partial_x^2 + \partial_y^2$ and $L \stackrel{\text{def}}{=} \partial_z(f_0/N)^2 \partial_z$, and LA is the back-rotated near-inertial velocity; the leading-order wave plus balanced flow velocity is

$$u + iv = LAe^{-if_0t} - \psi_y + i\psi_x, \quad (\text{A } 2)$$

where ψ is the streamfunction of the Lagrangian-mean (geostrophically balanced) flow. The PV is materially conserved,

$$q_t + J(\psi, q) = 0, \quad (\text{A } 3)$$

and the wave back-rotated velocity satisfies the YBJ equation,

$$LA_t + \frac{1}{2}f_0\Delta A + J(\psi, LA) + iLA\left(\frac{1}{2}\Delta\psi + \beta y\right) = 0. \quad (\text{A } 4)$$

The special family of solutions with a barotropic balanced flow $\psi = \psi(x, y, t)$, f -plane ($\beta = 0$), uniform background buoyancy frequency $N = N_0$, and $LA = e^{imz} \phi(x, y)$ wave velocity yields the vertical-plane-wave model in (2.7)-(2.8). The plane wave model is a solution of both XV and [Wagner & Young \(2016\)](#) equations because the barotropic flow assumption yields an infinite vertical-scale separation between waves and balanced flow.

A.2. Stokes drift

The horizontal component the Stokes drift of near-inertial waves is ([Wagner & Young 2016](#)):

$$u^S + iv^S = \frac{i}{f_0} (M_{zz}\partial^*M^* - M_z^*\partial^*M_z), \quad (\text{A } 5)$$

$$w^S = \frac{i}{f_0} (M_z^*\partial\partial^*M - \partial M_z\partial^*M) + \text{cc}. \quad (\text{A } 6)$$

where $M = (f_0/N)^2 A_z$ and $M_z = LA$, and we recall $\partial = (\partial_x - i\partial_y)/2$. In the vertical-plane-wave model, the back-rotated velocity is $M_z = \phi e^{imz}$, and the horizontal Stokes drift above simplifies to

$$(u^S, v^S) = (\mathcal{A}_y, -\mathcal{A}_x), \quad (\text{A } 7)$$

where $\mathcal{A} = |\phi|^2/2f_0$ is the action density. Thus the Stokes drift of the vertical-plane-wave model is horizontally non-divergent, with streamfunction $-\mathcal{A}$. The vertical component of the Stokes drift reduces to

$$\begin{aligned} w^S &= \frac{1}{f_0 m} [\partial(\phi\partial^*\phi^*) + \partial^*(\phi^*\partial\phi)] \\ &= \frac{1}{m} \left[\Delta \frac{1}{2} \mathcal{A} - \eta^{-1} \nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) \right], \end{aligned} \quad (\text{A } 8)$$

where we used (3.8) and the identities

$$\Delta = 4\partial\partial^* \quad \text{and} \quad J(f, g) = 2i(\partial^*f\partial g - \partial f\partial^*g). \quad (\text{A } 9)$$

An important property of the NIW-QG model is that there is no Lagrangian-mean vertical velocity:

$$w^E + w^S = 0. \quad (\text{A } 10)$$

Hence, using (A 8) and $\psi = \psi^E - \mathcal{A}$, we obtain an alternative expression for the potential vorticity (2.6) in terms of Eulerian-mean fields:

$$q = \Delta\psi^E + mw^E. \quad (\text{A } 11)$$

Appendix B. Quadratic conservation laws

B.1. Ehrenfest's theorem

To obtain (3.11) we begin by noting that with \mathcal{F} defined in (3.6)

$$\partial_t \mathcal{F} = \frac{i}{2} \lambda^2 (\phi_t \nabla \phi^* - \phi_t^* \nabla \phi) + \frac{i}{4} \lambda^2 \nabla (\phi \phi_t^* - \phi^* \phi_t). \quad (\text{B1})$$

Multiplying the wave equation by $i \nabla \phi^*$, adding to the complex conjugate, and using identities such as

$$\nabla \phi^* J(\psi, \phi) - \nabla \phi J(\psi, \phi^*) = J(\phi^*, \phi) \nabla \psi, \quad (\text{B2})$$

one eventually finds

$$\begin{aligned} \partial_t \mathcal{F} - \frac{i}{4} \lambda^2 \nabla (\phi \phi_t^* - \phi^* \phi_t) + \frac{1}{4} \lambda^2 \eta (\Delta \phi \nabla \phi^* + \Delta \phi^* \nabla \phi) = \\ - \nabla \psi \nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) + \eta \frac{1}{2} \zeta \nabla \mathcal{A} + \frac{i}{2} \lambda^2 (D_\phi \nabla \phi^* - D_{\phi^*} \nabla \phi). \end{aligned} \quad (\text{B3})$$

Taking the domain average, noting that the second and third terms on the left have zero average, and using the identity

$$\langle \nabla \psi \nabla \cdot (\hat{\mathbf{k}} \times \mathcal{F}) \rangle = \langle (\mathcal{F} \cdot \nabla) \hat{\mathbf{k}} \times \nabla \psi \rangle - \langle \hat{\mathbf{k}} \times (\zeta \mathcal{F}) \rangle, \quad (\text{B4})$$

we obtain (3.11) with the dissipative term

$$\varepsilon_{\mathcal{F}} \stackrel{\text{def}}{=} \frac{i}{2} \lambda^2 \langle D_\phi \nabla \phi^* - D_{\phi^*} \nabla \phi \rangle. \quad (\text{B5})$$

B.2. Wave potential energy

To obtain the wave potential energy equation (3.12) we take the dot product of $\nabla \phi^*$ with gradient of the wave equation (2.8) and add the complex conjugate; the calculation is best done using index notation. The final result is

$$\begin{aligned} \mathcal{P}_t + \nabla \cdot \left[\mathbf{u}^g \mathcal{P} + \frac{1}{2} \zeta \mathcal{F} + \frac{\lambda^2}{4} \frac{i}{2} \eta ((\nabla \phi \cdot \nabla) \nabla \phi^* - (\nabla \phi^* \cdot \nabla) \nabla \phi) \right] \\ = \frac{1}{2} \zeta \nabla \cdot \mathcal{F} - \frac{\lambda^2}{2} \phi_{,k} \sigma_{kl} \phi_{,l}^* + \frac{\lambda^2}{4} (\nabla \phi \cdot \nabla D_{\phi^*} + \nabla \phi \cdot \nabla D_{\phi^*}), \end{aligned} \quad (\text{B6})$$

where $\mathbf{u}^g \stackrel{\text{def}}{=} \hat{\mathbf{k}} \times \nabla \psi$ is the geostrophic velocity and

$$\sigma_{kl} \stackrel{\text{def}}{=} \frac{1}{2} (u_{k,l}^g + u_{l,k}^g) \quad (\text{B7})$$

is the geostrophic strain tensor. The local equation (B6) integrates to (3.12), with the dissipative term

$$\varepsilon_{\mathcal{P}} = \frac{\lambda^2}{4} \langle \nabla \phi \cdot \nabla D_{\phi^*} + \nabla \phi^* \cdot \nabla D_\phi \rangle = -\frac{\lambda^2}{4} \langle \Delta \phi^* D_\phi + \Delta \phi D_{\phi^*} \rangle. \quad (\text{B8})$$

B.3. Balanced kinetic energy

To obtain the balanced kinetic energy equation we first multiply the PV equation (2.7) by $-\psi$:

$$\mathcal{K}_t + \nabla \cdot [-\psi (\nabla \psi_t + \mathbf{u}^g q)] = \psi q_t^w - \psi \mathcal{D}_q. \quad (\text{B9})$$

To attack ψq_t^w on the right, we use the expression for q^w in (3.9). Thus

$$\psi q_t^w = \nabla \cdot \underbrace{\frac{1}{2} \left[\psi \left(\nabla \mathcal{A}_t + \frac{2}{\eta} \hat{\mathbf{k}} \times \mathcal{F}_t \right) - \nabla \psi \mathcal{A}_t \right]}_{\stackrel{\text{def}}{=} -\mathcal{K}_1} + \frac{1}{2} \zeta \mathcal{A}_t + \eta^{-1} \mathbf{u}^g \cdot \mathcal{F}_t. \quad (\text{B10})$$

Taking the dot product of (B3) with \mathbf{u}^g we have

$$\begin{aligned} \mathbf{u}^g \cdot \mathcal{F}_t = & \nabla \cdot \underbrace{\left\{ \mathbf{u}^g \left[\frac{1}{2} \lambda^2 (\phi \phi_t^* - \phi^* \phi_t) + \frac{1}{4} \eta \lambda^2 |\nabla \phi|^2 \right] - \frac{1}{4} \eta \lambda^2 [\nabla \phi \mathbf{u}^g \cdot \nabla \phi^* + \nabla \phi^* \mathbf{u}^g \cdot \nabla \phi] \right\}}_{\stackrel{\text{def}}{=} -\eta \mathcal{H}_2} \\ & + \eta \frac{1}{2} \zeta J(\psi, \mathcal{A}) + \frac{1}{2} \eta \lambda^2 \phi_{,k} \sigma_{kl} \phi_{,l}^* + \mathbf{u}^g \cdot \frac{1}{2} \lambda^2 (D_\phi \nabla \phi^* - D_{\phi^*} \nabla \phi). \end{aligned} \quad (\text{B 11})$$

Thus

$$\partial_t \mathcal{K} + \nabla \cdot [-\psi (\nabla \psi_t + \mathbf{u}^g q) + \mathcal{H}_1 + \mathcal{H}_2] = -\frac{1}{2} \zeta \nabla \cdot \mathcal{F} + \frac{1}{2} \lambda^2 \phi_{,k} \sigma_{kl} \phi_{,l}^* + \xi - \psi D_q, \quad (\text{B 12})$$

where

$$\xi = \frac{1}{2} f_0^{-1} (\phi^* D_\phi + \phi D_{\phi^*}) \frac{1}{2} \zeta + f_0^{-1} \mathbf{u}^g \cdot \frac{1}{2} (D_\phi \nabla \phi^* - D_{\phi^*} \nabla \phi) \quad (\text{B 13})$$

is the contribution of wave dissipation to the local balanced kinetic energy budget. Interestingly, the first term on the right of (B13) reveals that the dissipation of wave action in anti-cyclones is a source of balanced kinetic energy. The second term on the right of (B13) shows that the alignment of the ‘‘action-flux dissipation vector’’ $i(D_\phi \nabla \phi^* - D_{\phi^*} \nabla \phi)$, with the geostrophic velocity is also a source of balanced kinetic energy. The local equation (B12) integrates to the balanced kinetic energy equation (3.13) with the dissipative terms

$$\Xi \stackrel{\text{def}}{=} \langle \xi \rangle \quad \text{and} \quad \varepsilon_{\mathcal{K}} = -\langle \psi D_q \rangle. \quad (\text{B 14})$$

B.4. Specific expressions with biharmonic dissipation

The dissipative terms in (2.7) and (2.8) add small dissipation to the energy equations in section 3. The wave kinetic energy dissipation added to (3.1) is

$$\varepsilon_{\mathcal{K}} = -\nu_w \langle |\Delta \phi|^2 \rangle. \quad (\text{B 15})$$

The dissipation of wave potential energy in (3.12) is

$$\varepsilon_{\mathcal{P}} = -\frac{1}{2} \lambda^2 \nu_w \langle |\nabla \Delta \phi|^2 \rangle. \quad (\text{B 16})$$

Similarly, the balanced kinetic energy dissipation in (3.13) is

$$\varepsilon_{\mathcal{K}} = \kappa_e \langle \psi \Delta^2 q \rangle = \kappa_e \langle q \Delta^2 \psi \rangle. \quad (\text{B 17})$$

The wave dissipation contribution to the balanced kinetic energy budget is

$$\Xi = \frac{1}{2} \nu_w f_0^{-1} \langle \frac{1}{2} \zeta (\phi^* \Delta^2 \phi + \phi \Delta^2 \phi^*) \rangle + \nu_w f_0^{-1} \langle \frac{1}{2} \psi [J(\phi^*, \Delta^2 \phi) - J(\phi, \Delta^2 \phi^*)] \rangle. \quad (\text{B 18})$$

In all solutions of initial value problems reported in this paper, the dissipative terms (B15), (B16), (B17), and (B13) account for less—typically much less—than 10% of the energy tendencies.

REFERENCES

- BALMFORTH, N.J., LLEWELLYN SMITH, S.G. & YOUNG, W.R. 1998 Enhanced dispersion of near-inertial waves in an idealized geostrophic flow. *Journal of Marine Research* **56** (1), 1–40.
- BALMFORTH, N.J. & YOUNG, W.R. 1999 Radiative damping of near-inertial oscillations in the mixed layer. *Journal of Marine Research* **57** (4), 561–584.
- BARKAN, R., WINTERS, K.B. & MCWILLIAMS, J.C. 2016 Stimulated imbalance and the enhancement of eddy kinetic energy dissipation by internal waves. *Journal of Physical Oceanography* **47**, 181–198.

- BRETHERTON, F.P. & GARRETT, C.J.R. 1968 Wavetrains in inhomogeneous moving media. In *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, , vol. 302, pp. 529–554. The Royal Society.
- BÜHLER, O. & MCINTYRE, M.E. 1998 On non-dissipative wave–mean interactions in the atmosphere or oceans. *Journal of Fluid Mechanics* **354**, 301–343.
- BÜHLER, O. & MCINTYRE, M.E. 2005 Wave capture and wave–vortex duality. *Journal of Fluid Mechanics* **534**, 67–95.
- COX, S.M. & MATTHEWS, P.C. 2002 Exponential time differencing for stiff systems. *Journal of Computational Physics* **176** (2), 430–455.
- DANIOUX, E., VANNESTE, J. & BÜHLER, O. 2015 On the concentration of near-inertial waves in anticyclones. *Journal of Fluid Mechanics* **773**, R2.
- DANIOUX, E., VANNESTE, J., KLEIN, P. & SASAKI, H. 2012 Spontaneous inertia-gravity-wave generation by surface-intensified turbulence. *Journal of Fluid Mechanics* **699**, 153–157.
- FORNBERG, B. 1977 A numerical study of 2-d turbulence. *Journal of Computational Physics* **25** (1), 1–31.
- GERTZ, A. & STRAUB, D.N. 2009 Near-inertial oscillations and the damping of midlatitude gyres: a modeling study. *Journal of Physical Oceanography* **39** (9), 2338–2350.
- GRIMSHAW, R 1975 Nonlinear internal gravity waves in a rotating fluid. *Journal of Fluid Mechanics* **71** (3), 497–512.
- HOSKINS, BRIAN J 1975 The geostrophic momentum approximation and the semi-geostrophic equations. *Journal of the Atmospheric Sciences* **32** (2), 233–242.
- KASSAM, A-K & TREFETHEN, LN 2005 Fourth-order time-stepping for stiff pdes. *SIAM Journal on Scientific Computing* **26** (4), 1214–1233.
- KLEIN, P, LLEWELLYN SMITH, S.G. & LAPEYRE, G 2004 Organization of near-inertial energy by an eddy field. *Quarterly Journal of the Royal Meteorological Society* **130** (598), 1153–1166.
- LANDAU, L.D. & LIFSHITZ, E.M. 2013 *Quantum Mechanics: Non-relativistic Theory*, , vol. 3. Elsevier.
- MCINTYRE, MICHAEL E 2009 Spontaneous imbalance and hybrid vortex–gravity structures. *Journal of the Atmospheric Sciences* **66** (5), 1315–1326.
- MCWILLIAMS, J.C. 1984 The emergence of isolated coherent vortices in turbulent flow. *Journal of Fluid Mechanics* **146**, 21–43.
- MELESHKO, V.V. & VAN HEIJST, G.J.F. 1994 On Chaplygin’s investigations of two-dimensional vortex structures in an inviscid fluid. *Journal of Fluid Mechanics* **272**, 157–182.
- MOEHLIS, J. & LLEWELLYN SMITH, S.G. 2001 Radiation of mixed layer near-inertial oscillations into the ocean interior. *Journal of Physical Oceanography* **31** (6), 1550–1560.
- MURAKI, DAVID J, SNYDER, CHRIS & ROTUNNO, RICHARD 1999 The next-order corrections to quasigeostrophic theory. *Journal of the atmospheric sciences* **56** (11), 1547–1560.
- NAGAI, T., TANDON, A., KUNZE, E. & MAHADEVAN, A. 2015 Spontaneous generation of near-inertial waves by the Kuroshio Front. *Journal of Physical Oceanography* **45** (9), 2381–2406.
- SALMON, R 2016 Variational treatment of inertia-gravity waves interacting with a quasigeostrophic mean flow. *Journal of Fluid Mechanics* **809**, 502–529.
- SHAKESPEARE, C.J. & HOGG, A.McC. 2017 Spontaneous surface generation and interior amplification of internal waves in a regional-scale ocean model. *Journal of Physical Oceanography* .
- TAYLOR, S. & STRAUB, D.N. 2016 Forced near-inertial motion and dissipation of low-frequency kinetic energy in a wind-driven channel flow. *Journal of Physical Oceanography* **46** (1), 79–93.
- THOMAS, L. N. 2012 On the effects of frontogenetic strain on symmetric instability and inertigravity waves. *Journal of Fluid Mechanics* **711**, 620640.
- VANNESTE, J. 2013 Balance and spontaneous wave generation in geophysical flows. *Annual Reviews of Fluid Mechanics* .
- WAGNER, G.L. & YOUNG, W.R. 2015 Available potential vorticity and wave-averaged quasi-geostrophic flow. *Journal of Fluid Mechanics* **785**, 401–424.
- WAGNER, G.L. & YOUNG, W.R. 2016 A three-component model for the coupled evolution of near-inertial waves, quasi-geostrophic flow and the near-inertial second harmonic. *Journal of Fluid Mechanics* **802**, 806–837.

- XIE, J.-H. & VANNESTE, J. 2015 A generalised-lagrangian-mean model of the interactions between near-inertial waves and mean flow. *Journal of Fluid Mechanics* **774**, 143–169.
- YOUNG, W.R. & BEN JELLOUL, M. 1997 Propagation of near-inertial oscillations through a geostrophic flow. *Journal of Marine Research* **55** (4), 735–766.
- YOUNG, W.R., RHINES, P.B. & GARRETT, C.J.R. 1982 Shear-flow dispersion, internal waves and horizontal mixing in the ocean. *Journal of Physical Oceanography* **12** (6), 515–527.